

Interpersonal Psychotherapy: Healing with a Relational Focus

Abstract: Interpersonal psychotherapy (IPT) is a time-limited psychotherapy that focuses on relationship stressors and ways to adaptively engage with social supports. Since the first controlled IPT depression study 40 years ago, new applications of the model have emerged, informed by research and public health needs. Evidence for its effectiveness has led to its inclusion in expert consensus treatment guidelines for the treatment of depression, eating disorders, and bipolar disorder. This paper provides a clinical synthesis of IPT, reviewing adaptations that include: IPT-A, for use with adolescents with depression; interpersonal social rhythm therapy (IPSRT), for patients with bipolar disorder; IPT, for patients with eating disorders; and IPT, for patients with depression in culturally diverse settings. With its clear clinical guidelines, therapist- and patient-friendly approach, and data supporting its effectiveness, IPT is easily integrated into mental health care to help patients with mood or eating disorders and interpersonal problems.

Relationships matter—in health, disease, coping with stress, and recovering from illness. This is the rationale for interpersonal psychotherapy (IPT) (1), a time-limited psychotherapeutic model that focuses on relational aspects of experience and mental health. IPT treats depression across the life span (2–4), and has been successfully adapted for differing clinical populations, including those with bipolar disorder (5), posttraumatic stress disorder (6, 7), and eating disorders (8, 9). IPT can be effectively delivered by a variety of health providers from mental health specialties such as psychiatrists, psychologists, nurses, social workers, and occupational therapists; to trained lay health workers (10). This range of treatment providers can help to address the wide global gap between the need for, and access to, mental health care (11, 12).

The first controlled IPT study for depression was published 40 years ago (13). Since then, novel applications of the model have emerged, informed by research and public health needs. Evidence for its antidepressant effects, established through numerous clinical trials (14) has led to its inclusion in expert depression treatment guidelines (Table 1) (16, 17, 24) including those of the World Health Organization (WHO) (18). Consensus guidelines for the treatment of eating disorders and bipolar disorder also recommend IPT, based on level 1 evidence of at least two randomized clinical trials conducted by two separate groups of investigators demonstrating positive outcomes (20–22).

At the time of IPT's genesis, Bowlby's seminal work on attachment theory (25, 26), Brown and Harris' studies on the associations between bereavement and depression (27), and the etiological links between biological and psychosocial factors were becoming influential in the discourse on illness and recovery (28, 29). Since that time the importance of relationships for health, coping, and resilience has become well established (30–33). Mental illnesses are often triggered or exacerbated by relationship stressors, such as interpersonal losses, life changes, loneliness, or conflicts. IPT treatment provides therapeutic roadmaps to work through these core relationship focus areas.

Author Information and Disclosure

Paula Ravitz, M.D., F.R.C.P.C., Department of Psychiatry, University of Toronto, and Department of Psychiatry, Mount Sinai Hospital, Toronto.

Priya Watson, M.D., F.R.C.P.C., Department of Psychiatry, University of Toronto, and Centre for Addiction & Mental Health (CAMH), Toronto.

The authors receive book royalties and report no other financial relationships with commercial interests.

The authors thank Jamie Park in the Mt. Sinai Hospital, Department of Psychiatry Research Office for assistance with manuscript review and editing, the Mt. Sinai Hospital Morgan Firestone Psychotherapy Chair, the CAMH Office of Education, Dr. Alan Ravitz, Susan Biglieri, and Edward McAnanama for their clinical wisdom and assistance with the manuscript. No funding was received for this project.

Address correspondence to Dr. Ravitz, 600 University Ave., Room 930, Mount Sinai Hospital, Toronto, Ont., Canada, M5G 1X5; e-mail: pravit@mtsinai.on.ca

Table 1. Consensus Treatment Guidelines that Include IPT

Clinical Population	Experts' Consensus Group
Depression	National Institute of Clinical Excellence (NICE) (15) American Psychiatric Association (APA) (16) Canadian Network for Mood and Anxiety Treatments (CANMAT) (17) World Health Organization (WHO) (18) Royal Australian and New Zealand Clinical Practice Guidelines (RANZCP) (19)
Eating Disorders	APA (20)
Bipolar Disorder	APA (21) CANMAT (22) RANZCP (23)

This paper provides an overview of IPT (1) with an emphasis on the clinical guidelines as summarized in Figure 1 (34). It is beyond the scope of this clinical synthesis to describe all IPT adaptations in depth; however, we briefly review those included in consensus treatment guidelines (Table 1). These include: IPT-A, for use with adolescents with depression; IPSRT, for patients with bipolar disorder; IPT, for patients with eating disorders; and IPT, for patients with depression in culturally diverse settings. We begin by describing and discussing the evidence supporting each adaptation of IPT and then review the components of the original treatment.

IPT-A

Depression poses severe risks to adolescents, including incomplete schooling, substance abuse, early pregnancy, self-harming behaviors, recurrent depressive episodes, and physical morbidity. By age 18, an estimated 20% of adolescents will have had a major depressive episode (35). An intervention to treat, and potentially prevent the recurrence of depression, is therefore of great public health significance. Mufson and colleagues created and piloted the IPT-A model during the 1990s, when there were no published controlled trials for any individual psychotherapy for depressed adolescents (36). Since then, IPT-A has been established as an effective treatment in numerous controlled trials (2, 37, 38). IPT-A adheres to the core tenet that relationships are the locus of intervention and healing, but also contains important developmentally-oriented modifications. These modifications take into account the differences in life circumstances and maturation tasks that confront adolescents (39). Aligned with the de-

velopmental needs necessary for healthy adolescent maturation, areas of therapeutic work in IPT-A include: reducing conflict in relationships, developing and using healthy communication strategies, resolving identity crises, and building and maintaining a supportive set of relationships with the family and peers.

IPT-A remains a brief psychotherapy (12 sessions) with distinct phases and the same focal areas as IPT. It also employs strategies for engaging adolescents, such as telephone contact between sessions. Another important modification of IPT-A is including parents or other primary caregivers in all phases of the therapy process. Parents join the diagnostic assessment and psychoeducation at the beginning, including conveying a "limited sick role" to the adolescent. With the adolescent's permission, parents may attend some middle phase sessions, to orient them to the new strategies the teen will be trying at home, and where needed, to address disputes within the family. In the termination phase, both the adolescent and parents will join in a review and consolidation of gains, with discussion of early signs of relapse, and development of a plan for possible future depression recurrence.

VIGNETTE 1: AN EXAMPLE OF AN IPT-A CASE WITH A FOCUS ON DISPUTES

Greg is a 15-year-old adolescent with a 1-year history of low mood, anergia, anhedonia, and difficulty with concentration and academic performance. He denied suicidal ideation and substance abuse. At the time of his assessment, Greg rarely attended school, which was the subject of near-daily arguments with his mother. He had stopped all extracurricular activities, including soccer and drama, which he used to enjoy. He typically stayed awake for most of the night, often playing computer games online, and slept during the day. His absences from school resulted in failing grades and increasing social isolation from his friends.

Greg was an only child who lived with his mother. When questioned about his life at the time of the onset of his depressive symptoms, he recounted a series of verbal conflicts with his parents who had separated two years ago. Greg cut off contact with his father a year ago, feeling hurt by what he perceived as a lack of support and interest.

In IPT-A, meetings with a parent are held in the beginning phase, and in this case the therapist met Greg and his mother to discuss the diagnosis of depression, instill hope, and provide psychoeducation on depression, IPT, and the impact of depression on Greg's willingness and ability to succeed at school. They discussed that Greg's social isolation, frequent conflicts with his mother, and alienation from his

Figure 1. IPT for Depression Pocket Card.

PRACTICE REMINDER IPT for Depression

Beginning Phase Tasks


1. Conduct an assessment. Establish the therapeutic alliance.
2. Learn about problems associated with the onset of the symptoms and the people that help or hinder (with the interpersonal inventory).
3. Formulate the IPT focus: bereavement, life changes (role transitions), disagreements or interpersonal sensitivity. When choosing the focus, ensure a chronological link to the onset or perpetuation of symptoms; associated dysphoric affect; possibility for working through change.
4. Discuss goals—to remit symptoms, improve functioning, and work through agreed-upon problem area.
5. Provide psychoeducation: convey your understanding of the patient's suffering as disabling, not the patient's fault, connected with triggers, treatable, and that she or he needs social support during this time. IPT will help with current interpersonal problems that are linked to distress.

Middle Phase Sessions: When patients discuss their problems and begin to make changes in their lives. Link the focal area to the onset or perpetuation of symptoms and explore problems, relationships, and interactions associated with worsening or improvement of their distress. Focus on communication, recruiting supports, and ways to handle interpersonal problems. Track symptoms, safety, functioning, and progress weekly using validated screening questionnaires.

Concluding Phase Tasks: Review experience of treatment, changes (in symptoms and in relationships), successes, and efforts, strengths and supports. Discuss how to discern a "bad day" from the return of clinical depression, the signs of recurrence, and contingency planning.

Adapted & Reprinted from "Psychotherapy Essentials to Go: Interpersonal Psychotherapy for Depression" © 2013 by Paula Ravitz, Priya Watson, Sophie Grigoriadis. With permission of the publisher, W.W. Norton & Company, Inc.

- Goals:
- Reduce symptoms by improving social functioning
 - Helps patients to:
 - Repair or build supports
 - Improve communication
 - Resolve interpersonal problems



Focus-Specific Tasks

Life Changes—Social Role Transitions

1. Encourage the patient to describe in detail the change with its associated disruptions.
2. Explore positive aspects of the old role (mourning losses), and challenges and opportunities of the new role/life changes.
3. Identify options to improve the situation and people who can help to make the new role easier.

Bereavement

1. Explore in detail the circumstances of the loved one's death.
2. Encourage the patient to tell of his or her relationship with the deceased (how they met, how their relationship evolved, the positive aspects and the disappointments).
3. Throughout, encourage the patient to interact with caring friends and family and to find meaningful ways to use his or her time.

Disputes

1. Explore in detail the relationship, the client's view of the issues in dispute, what she has tried, and what she would like to change.
2. Explore her sense of the problem from the other person's point of view, how they impact one another, and her understanding of the other person's reactions and feelings.
3. Explore specific examples of recent exchanges (communication analysis) to clarify each party's position; explore differing values and expectations; and foster understanding of one another's perspectives and limitations, in order to come up with more effective choices and ways of communicating.

father might be driving his depression. The therapist explained that Major Depressive Disorder was a treatable condition and that Greg would feel and function much better as his depressive symptoms began to remit.

Middle phase sessions focused on the disputes with Greg's parents and included some joint sessions with his mother; however, he did not wish his father to come in. Greg and his mother collaboratively devised communication strategies for their arguments. After a number of sessions exploring wishes and expectations, doing role plays and problem-solving,

Greg's relationships with both parents improved, and he restored contact with his father. Over time, Greg contacted friends and began to re-engage in his social life. By 8 weeks of IPT-A, his depressive symptoms remitted and he was functioning well at home and at school. At the termination of therapy, Greg acknowledged, with relief, the significant improvements in his mood and close relationships, and his therapist emphasized that Greg now had a set of communication tools and relationship strategies that he could use whenever he experienced low mood.

INTERPERSONAL SOCIAL RHYTHM THERAPY (IPSRT)

Bipolar disorder is a chronic psychiatric illness characterized by alternating depressive and [hypo] manic mood episodes that can cause serious negative consequences to the patient and his or her family if left untreated. Rates of conversion from unipolar to bipolar disorder are relatively low. However, bipolar disorder is commonly misdiagnosed as unipolar depression; and treatment-refractory depression may, in some cases, represent occult bipolar disorder, with its attendant greater severity and functional impairment. Although bipolar disorder is much less prevalent than major depressive disorder, it is challenging to treat and often impacts upon the life circumstances and relationships of patients and their families in debilitating ways such that bridges are burned, and social development is derailed (5). Frank, Swartz, and colleagues created IPSRT for this patient population, based on the findings that radical changes in daily routines precede circadian rhythm instability, which can lead to affective episodes (40, 41). IPSRT utilizes the Social Rhythm Metric (SRM), a behavioral tool (42) developed to help stabilize circadian and social rhythms during manic or depressive phases of bipolar illness and to provide patients with skills that will help prevent the onset of future episodes (43, 44). The SRM tracks five distinct activities: the times they get out of bed; have first contact with another person; start meaningful activity such as work, school, housework or volunteering; have dinner; and go to bed. Patients also record whether they were alone or with others when they participated in each daily activity. Finally, patients rate their mood and energy level each day. Therapists use the SRM to identify links between fluctuations in mood and regularity of routines. Over time, unstable social rhythms that negatively impact mood or energy are identified as targets for change, moving toward a balance of rest, activity, and personal contacts that promote mood stability.

Frank et al. suggest combining mood stabilizing medication, IPT, and the simple SRM behavioral tool to restore circadian and social rhythms during the acute and maintenance phases of treatment, helping to stabilize symptoms and decrease the risk of relapse (45). Compared with intensive clinical management, patients with bipolar disorder treated with IPSRT have reduced rates of relapse and higher regularity of social rhythms at the end of acute treatment. Additionally, IPSRT is shown to be effective in reducing suicidal behavior in patients with bipolar disorder (46). As IPSRT has been shown to delay relapse, speed recovery, and increase both occupational and psychosocial functioning (43), it has been

recommended in consensus treatment guidelines for bipolar disorder (21).

IPT FOR EATING DISORDERS

Eating disorders such as bulimia nervosa, anorexia nervosa, and binge-eating disorder represent a public health concern due to their increasing prevalence (47) and association with elevated mortality rates (48). IPT has been studied with patients who have these eating disorders, with the target of treatment being the interpersonal difficulties that maintain symptoms (8, 9, 49). Randomized controlled trials have indicated that, for patients with bulimia nervosa or binge-eating disorder, IPT takes longer than Cognitive Behavioral Therapy (CBT) to achieve recovery; however, both models of treatment are equally effective.

Both models, despite having differing emphases—for CBT to modify distorted thinking about shape and weight, and for IPT to modify interpersonal functioning—have been shown to effect changes in both cognitions and relationships (49). Long-term recovery rates for patients with anorexia nervosa hover around 49%, whether treated with CBT or IPT, which suggests that supplementary or longer treatment approaches may be needed for anorexia nervosa (50).

IPT IN INTERNATIONAL AND TRANS-CULTURAL SETTINGS

Underdetected and untreated mental illness has been identified by the World Health Organization as a severe global problem (51), for which many global mental health initiatives are now underway. IPT has been at the forefront of efforts to increase access to culturally appropriate psychotherapy for common mental disorders. Adaptations of the IPT model with far-reaching public health scope have been developed for a number of international and trans-cultural settings. Over the past decade, IPT has undergone significant cultural and methodological reworking, for implementation in settings such as Uganda, Rwanda, India, and Ethiopia (52–58) and for use among specific cultural populations in North America (59, 60). These studies addressed the intersection of culture, illness, and healing. Significant changes were made to diagnostic instruments in these studies, with piloting and cultural validation of the language and concepts used (61, 62). Local health-care workers were trained to deliver IPT, increasing the capacity for ongoing local service provision once the studies were complete. Treatment groups were chosen to reflect the local reality and public health needs, rather than “pure” depression cases; such as the work done with internally displaced youth in

northern Uganda, who presented with ongoing severe psychosocial stressors and numerous psychiatric comorbidities (53).

These international IPT studies have contributed to the advancement of global psychotherapy research through their innovative methods. The Uganda IPT trials, for instance, began by developing mechanisms for the cross-cultural adaptation and validation of assessment instruments. This ensured that the diagnostic category of depression, which varies in meaning across cultures and social contexts, was a valid measurement category (54). These studies were also groundbreaking as they were the first published controlled clinical trials of psychological interventions in sub-Saharan Africa, and they provided an early evidence base for the feasibility of future psychotherapeutic interventions in this region (54, 58). Given the 2004 WHO findings that 76–85% of the individuals with serious mental illness who lived in less-developed countries received no treatment in the preceding year, studies that demonstrate the validity and feasibility of an effective treatment provide an essential public health service. These studies provide a rationale for resource allocation and systemic effort for treatment of these disorders. This is particularly important in resource-poor and underserved countries where the physical and mental health needs are great and the spending constraints are severe.

IPT FORMATS

IPT can be delivered in differing formats and doses. For group IPT, additional therapeutic processes to capitalize on the dynamic therapeutic elements of group, such as cohesion, universality, group learning, and receiving constructive feedback from other members, are integrated into the treatment (63). In addition, there are ultra-brief versions of IPT, one which focuses on screening, support, and triage (64), and another, called Interpersonal Counseling (IPC) that with as few as six sessions, using the same core principles as IPT, has been shown to be helpful in primary care settings (65). Having outlined the background and several variations of IPT, we will now describe the clinical guidelines that apply regardless of dose (4–16 sessions) or format (individual or group therapy).

IPT CLINICAL GUIDELINES

Like all time-limited psychotherapy treatments, IPT has beginning, middle, and termination phase tasks. The IPT guidelines must still be delivered with attention to good therapeutic processes such as using empathy, mentalizing, and reflection with positive

regard and respectful engagement to foster a strong therapeutic alliance.

BEGINNING PHASE OF IPT

The beginning phase, akin to all psychiatric and psychotherapeutic treatments, entails getting to know your patient, establishing a therapeutic alliance, and clarifying the goals that will guide the treatment. A psychiatric assessment is done to learn about the individual patient, his or her suffering, symptoms, and important close relationships. We suggest adding cultural formulation questions (66) to learn about the meaning of symptoms or illness, current practices of coping, and expectations for care. Explicitly inquiring about the cultural aspects of the presentation will serve to increase the therapist's understanding of the patient, reduce misrecognition of symptoms, and likely improve the therapeutic alliance (67).

An important goal of the initial phase of IPT is establishing the interpersonal problem area. In IPT's standard formulation, the therapist chooses among four possible problem areas: 1) grief (difficulty mourning the death of an important person in the patient's life), 2) role transitions (change in major social roles such as graduation, divorce, retirement, and job promotion), 3) role disputes (nonreciprocal role expectations between the patient and a significant other), or 4) interpersonal sensitivities (long-standing impoverished or contentious relationships).

If symptoms and functional impairment are high or chronic, then adjunctive antidepressant pharmacotherapy should be considered and discussed with the patient. The beginning phase involves gathering and communicating much information, including delivering psychoeducation about the mental disorder and its treatment, the rationale and description of IPT, and the instillation of hope for recovery.

During the beginning phase, the therapist gathers an expanded psychosocial history called the "Interpersonal Inventory." The Interpersonal Inventory consists of a careful review of the people in the patient's current relationships and life circumstances who are close, or with whom there are distressing conflicts. This is done in order to identify those who help or hinder in times of need. In the process of conducting the Interpersonal Inventory, a clinician gains an appreciation of a patient's significant past and present relationships. Losses, tensions, disagreements, trauma, and patterns of relating are revealed through the Inventory, helping to highlight relationships that may be appropriate for therapeutic focus. For example, a patient might become tearful, or less coherent with a reduced capacity to be clearly descriptive of specific relationships during

the Inventory process, signaling possible unresolved trauma. The beginning phase ends when an IPT focus is collaboratively chosen for the middle phase. This requires transparent discussion between the therapist and the patient about which interpersonal focal area and which relationships will comprise the focus of treatment. Getting agreement on the IPT focus sets the stage for doing work on the interpersonal problem area most saliently linked to the current illness episode. Although a relatively discrete and specific interpersonal issue is chosen as the focus of IPT treatment (see below), the IPT problem area is often the proverbial tip of the iceberg. The problem area may be indicative of more pervasive interpersonal difficulties such as chronic relationship disputes or maladaptive patterns of coping with stress. The time-limited intervention of IPT does not profess to bring about significant character change; however, the discoveries and changes made over the course of therapy, although brief, can be generalized beyond current circumstances.

In the middle phase, the IPT therapist facilitates reflection, making links between distress or symptoms, relational experiences, and the interpersonal problem area. The early middle phase involves fostering a deepened understanding of current relationships, interpersonal problems, and experiences; whereas in the latter middle phase there is a push for change as patients generate ideas for problem solving, interpersonal activation, and engagement with supports.

MIDDLE-PHASE FOCUS SPECIFIC THERAPEUTIC GUIDELINES

GRIEF

In the case of bereavement, during the early middle phase, the details of the death, burial, and acute period of grieving are reviewed. The guidelines provided by Klerman, Weissman, and Markowitz provide a powerfully helpful therapeutic roadmap for therapists to guide patients through the process of working through emotionally painful losses (1). The therapist guides the patient to reflect on the lost significant other and this relationship, returning with greater depth of exploration than when first described during the Interpersonal Inventory. For relationships in which there were ambivalent feelings and conflict, it is important to pace the exploration slowly, so that there can be emotional processing to integrate traumatic aspects of the loss. In the latter middle phase of working with grief, ways to “move on,” and to replace aspects of what has been lost from this relationship through other social supports are discussed.

ROLE TRANSITIONS

The notion of *social roles* provides a helpful conceptual idea for the multiplicity of expectations in relationships that are held in differing contexts and with differing levels of intimacy. These include relationships within families, romantic connections at work or school, and in community or religious groupings. Social roles define expectations in relationships, and inevitably change over time. Life changes, with their accompanying social role changes, are called “role transitions” in IPT. Role transitions are situated both externally and internally—in their impacts upon relationships, and the patient’s sense of self. Transitions can be planned or unplanned, wished for or dreaded. Examples of role transitions include: becoming a new parent; marrying or divorcing; migrating; becoming medically or mentally ill; and changes in vocational status such as becoming promoted, unemployed, or retired. Role transitions, with their demands and stresses, can precipitate or worsen states of mental illness such as depression. IPT techniques for addressing role transitions include exploring the change with its losses, challenges, and opportunities. For instance, there may be positive aspects of an “old role” that could be carried forward in some way. Opportunities presented by the new role include changes in appreciation, perspective, relationships, and behaviors to better align with a patient’s values, sense of well-being, and meaningful engagement.

VIGNETTE 2

The next example is of an IPT case with a focus on role transitions. Patricia is a 37-year-old married mother of a healthy 4-week-old baby boy. She was referred by her family physician for IPT treatment of postpartum depression. This was a planned pregnancy and there were no complications. Patricia worked as an accountant but was currently on maternity leave. Although she had been excited to become a new mother, she was surprised to find herself tearful and sad most days, with a diminished sense of self-esteem. Though she was highly accomplished as an accountant, she had trouble breastfeeding and felt incompetent, out of control, and overwhelmed as a new mother. She also felt guilty that she experienced little pleasure or connection with her newborn son. She was not suicidal and had no thoughts of harm to her child.

The Interpersonal Inventory, conducted in session 2 after the assessment, revealed that she felt estranged from her spouse and, unaccustomed to asking for help from others, felt quite isolated. Prior to becoming a mother, both she and her husband worked very long hours and socialized mainly with their single friends who were not parents.

In keeping with the middle phase tasks of IPT for role transitions, the therapist explored how Patricia's life had changed, her sense of having lost the positive aspects of her life before the birth, when she had more flexibility and a lessened sense of responsibility at home. She also voiced confusion about managing her new social role, feeling overloaded with responsibilities and the need to master child care, breastfeeding, and emotionally connecting with her baby. Patricia worked on giving herself permission to gradually learn the many new skills she needed, including breastfeeding, and to recruit support from her extended family, who were happy to help. Over the latter middle phase, as her depression remitted, she became more comfortable asking for help from others, especially her spouse. She came to embrace and enjoy the opportunities of her new role and of her relationships with her spouse and their son.

ROLE DISPUTES

There is a sequence of therapeutic tasks in the focal area of disputes. The first of these is to engage, through detailed exploration, with the relationship with the disputed other. The core issues in the dispute, such as transgressions of trust, disparate values, or nonshared expectations, are identified, with a focus on promoting reciprocal understanding. This goes hand-in-hand with improving communication via unpacking upsetting interactions (see Communication Analysis below) and behavioral experiments of interacting from a place of better understanding. Interpersonal expectations are explored and sometimes adjusted toward what is both reasonable and realistic, considering the capacity and perspectives of the other. By working on a current dispute, problematic patterns of relating in which a patient inadvertently "fuels the fire" of the conflict, or creates interpersonal distance, can be helpfully addressed and may often be generalized to improve relationships more broadly.

Interpersonal Sensitivity, also known as *deficits* in earlier descriptions of IPT, is a problem area that is sometimes left out of adaptations of the manual. Considered a default category by some, this focus is chosen when there is an absence of life events or close relationships. In cultural adaptations of IPT used in low and middle income countries with communal societies, deficits in social supports are not usually a focus of treatment; however, recruiting of social supports is a pan-focal task used with all IPT patients. In process research, this focal area has not been implicated as having worse outcomes (68).

For the focal area of interpersonal sensitivity, rather than striving for full "resolution" of the problem area, the goal is to "lessen" it. An IPT strategy for interpersonal deficits is to review negative and positive

aspects of past significant relationships (e.g., friendships, romantic relationships, and even past therapeutic relationships). This can reveal patterns of relating that are interpersonally distancing, and that may be recapitulated in the current therapeutic relationship. Unlike other IPT focal areas, the therapeutic relationship can be psychodynamically explored in conjunction with an active focus on expanding social supports.

VIGNETTE 3

A 41-year-old male consultant who presented with chronic depression that had recently worsened in the context of being rejected by an online dating site described having no close relationships. During sessions, he was averse to expressing or identifying emotions and was quite sophisticated in drawing the therapist into intellectual themes. He spoke with frustration of his past experience of psychotherapy in which he felt unhelped, and said that his therapist "didn't get to how I felt." This statement provided leverage to collaboratively explore his emotional experience of the current therapy session. The patient identified his use of his intellect to engage with people, yet found the friendships distant and unsatisfactory. In becoming more expressive of his inner, emotional experience of relationships in the here-and-now of the session, the patient felt understood and helped. Roles plays, communication analyses, and brainstorming were used to help improve his interactions with others. Over the remaining course of IPT sessions, his depressive symptoms lessened, he became more in touch with his own emotions, and as well, he became involved in a community-based project, volunteering his skills in a hobby he enjoyed that decreased his social isolation.

In the case of social deficits, as illustrated in the case above, brainstorming with the patient on ways to reduce social isolation and engage with others who share the patient's interests or values can help to alleviate depressive symptoms. Empathic exploration of internal and external experiences, such as emotions, thoughts, reactions, behaviors, communication, and interactions, is combined with judicious exploration of interpersonal dynamics that may emerge in the therapeutic relationship itself. "Plain old therapy" techniques as elaborated in psychodynamic (69) and mentalizing-based treatments (70, 71) can be helpfully integrated within the time-limited structure of IPT for patients with social deficits, comorbid personality disorders, and interpersonal sensitivity (72).

OTHER IPT THERAPEUTIC STRATEGIES

Communication Analysis is a helpful therapeutic strategy used in all focal areas to explore distressing

interactions that are associated with worsening symptoms. By unpacking specific examples of conversations and exploring an interaction, communication analysis creates the space to reflect and identify problems such as misunderstandings, communicating in ways that inadvertently fuel tensions, or having a lack of empathy. Using role plays or exploratory questions, such as asking a patient, “What would you like this person to understand?” can generate insight and ideas for change. In the process of communication analysis, a patient can discover and express more clearly his or her feelings and expectations. These can then be validated by the therapist, further explored, and sometimes revised. In this way, communication analysis is a collaborative means to improving communication behaviors, not just of recollecting troubling interactions.

RECRUITING AND UTILIZING SOCIAL SUPPORTS

Regardless of the focal area, IPT promotes patients’ recruiting and utilizing helpful psychosocial supports, such as family members, friends, colleagues, or community members. The goal is to help patients to gain comfort, understanding, and help from others, as the association of interpersonal support and psychological well-being has been well established (73).

TERMINATION PHASE OF IPT

The IPT *concluding phase guidelines* can be generalized to many therapeutic contexts (e.g., outpatient clinic or discharge from an inpatient or day hospital setting) and can provide a helpful set of therapeutic actions to consolidate gains and plan for the event of possible relapse. In the final 1–2 sessions, the therapist invites the patient to reflect on his or her experience of treatment and what he or she is taking away from the therapy. This reflection might include reviewing strategies to understand and cope with future stressors and use social supports, and perspectives on how the patient or his or her situation may have changed. Early signs of relapse are reviewed, and a contingency plan for intervention is created. By explicitly attending to this process of “saying goodbye” in treatment, acknowledging the work a patient has done and feelings or worries about ending therapy, the termination phase tasks provide therapeutic opportunities to empathically support the patient’s self-efficacy and improved coping strategies.

As with all therapeutic treatments, it should be noted that IPT does not work for all patients. Research about “what works best for whom” (74), and meta-analyses on the effects of therapist, patient, therapeutic relationship, and treatment factors on outcomes (75) highlight the importance of psycho-

therapeutic common factors, especially the therapeutic alliance. IPT process research has begun to examine what mitigates the response to treatment, including factors such as motivation, chronicity, attachment patterns of relating, and interpersonal problems (76–83).

To improve IPT outcomes, simple modifications that attend to individual patient differences have been recommended. For patients with panic symptoms, these modifications include focusing on emotions, especially fear, and somatic symptoms as cues of interpersonal distress. For patients with chronic or severe depression symptoms, the course of treatment may be extended, and medication concurrently used. Other modifications include an expanded biopsychosocial etiological case formulation, to include attachment style, culture, and self-definition (84).

Common factors exist among all effective therapies, related to the therapeutic alliance with empathy, rapport, positive regard, genuineness, and responsiveness (85). However, there are differences between psychotherapies—in their goals, frame, and techniques. Cognitive Behavior Therapy, also time-limited with goals of remitting symptoms of mental illnesses such as depression, differs from IPT in its focus on the links among thoughts, emotions, and behaviors with a high level of structure, and assigned between-session homework, for example, using automatic thought records, behavioral activation, and graded exposure (86). Psychodynamic Therapy is usually open-ended in its time frame, and informed by psychoanalytic theories that privilege how the past can influence our experience of the present. During sessions, rather than choosing a specific focal area, the therapist focuses on affect and expression of feelings, identifying distressing or self-defeating patterns, with exploration of associations, dreams, and transference dynamics (how we may project expectations from past relationships into present ones including within the therapeutic alliance). The goals of psychodynamic therapy include personality change and improved intimacy in relationships (87, 88). Supportive Therapy differs from IPT in that it does not have a specific focus on relationships or interpersonal problems, but rather on coping, building self-esteem, and reducing anxiety (89).

In conclusion, IPT is a powerfully helpful time-limited treatment for depression, across the lifespan and in differing cultures. It provides a clear structure with phase and focus-specific guidelines. IPT has been successfully used to treat depression, bulimia, and binge-eating disorders, and as an adjunctive treatment for bipolar disorder. There is growing evidence for its effectiveness in addressing other psychiatric problems, such as posttraumatic stress, and its use in differing formats, such as by

telephone or Internet. Even with complex and chronically ill patients, such as those frequently seen in community-based “real world” settings, IPT’s relationally-focused principles and guidelines are easily integrated and modified to improve outcomes in mental health care.

REFERENCES

- Weissman M, Markowitz JC, Klerman GL: *Clinician’s Quick Guide to Interpersonal Psychotherapy*. New York, Oxford University Press, 2007
- Mufson L, Dorta KP, Wickramaratne P, Nomura Y, Olsson M, Weissman MM: A randomized effectiveness trial of interpersonal psychotherapy for depressed adolescents. *Arch Gen Psychiatry* 2004; 61:577–584
- Reynolds CF 3rd, Dew MA, Pollock BG, Mulsant BH, Frank E, Miller MD, Houck PR, Mazumdar S, Butters MA, Stack JA, Schlermitzauer MA, Whyte EM, Gildengers A, Karp J, Lenze E, Szanto K, Bensasi S, Kupfer DJ: Maintenance treatment of major depression in old age. *N Engl J Med* 2006; 354: 1130–1138
- Reynolds CF 3rd, Frank E, Dew MA, Houck PR, Miller M, Mazumdar S, Perel JM, Kupfer DJ: Treatment of 70(+)-year-olds with recurrent major depression; excellent short-term but brittle long-term response. *Am J Geriatr Psychiatry* 1999; 7:64–69
- Frank E, Swartz HA, Kupfer DJ: Interpersonal and social rhythm therapy: managing the chaos of bipolar disorder. *Biol Psychiatry* 2000; 48:593–604
- Markowitz JC, Milrod B, Bleiberg K, Marshall RD: Interpersonal factors in understanding and treating posttraumatic stress disorder. *J Psychiatr Pract* 2009; 15:133–140
- Talbot NL, Chaudron LH, Ward EA, Duberstein PR, Conwell Y, O’Hara MW, Tu X, Lu N, He H, Stuart S: A randomized effectiveness trial of interpersonal psychotherapy for depressed women with sexual abuse histories. *Psychiatr Serv* 2011; 62:374–380
- Fairburn CG, Jones R, Peveler RC, Hope RA, O’Connor M: Psychotherapy and bulimia nervosa; longer-term effects of interpersonal psychotherapy, behavior therapy, and cognitive behavior therapy. *Arch Gen Psychiatry* 1993; 50:419–428
- Wilson GT, Wilfley DE, Agras WS, Bryson SW: Psychological treatments of binge eating disorder. *Arch Gen Psychiatry* 2010; 67:94–101
- Patel V, Weiss HA, Chowdhary N, Naik S, Pednekar S, Chatterjee S, De Silva MJ, Bhat B, Araya R, King M, Simon G, Verdelli H, Kirkwood BR: Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. *Lancet* 2010; 376:2086–2095
- Belkin GS, Unuetzer J, Kessler RC, Verdelli H, Raviola GJ, Sachs K, Oswald C, Eustache E: Scaling up for the “bottom billion”: “5x5” implementation of community mental health care in low-income regions. *Psychiatr Svcs* 2011; 62:1494–1502
- van Ginneken N, Tharyan P, Lewin S, Rao GN, Meera SM, Pian J, Chandrashekar S, Patel V: Non-specialist health worker interventions for the care of mental, neurological and substance-abuse disorders in low- and middle-income countries. *Cochrane Database Syst Rev* 2013; 11:CD009149
- Klerman GL, Dimascio A, Weissman M, Prusoff B, Paykel ES: Treatment of depression by drugs and psychotherapy. *Am J Psychiatry* 1974; 131:186–191
- Cuijpers P, Geraedts AS, van Oppen P, Andersson G, Markowitz JC, van Straten A: Interpersonal psychotherapy for depression: a meta-analysis. *Am J Psychiatry* 2011; 168:581–592
- Depression: the treatment and management of depression in adults (update). National Institute for Health and Care Excellence; 2010 [cited 2014 Feb 21]; Available from: <http://guidance.nice.org.uk/CG90/NICEGuidance/pdf/English>.
- American Psychiatric Association: *Practice Guideline for the Treatment of Patients with Major Depressive Disorder*, 3rd ed. Arlington, VA, American Psychiatric Association, Inc, 2010
- Parikh SV, Segal ZV, Grigoriadis S, Ravindran AV, Kennedy SH, Lam RW, Patten SB; Canadian Network for Mood and Anxiety Treatments (CANMAT): Canadian Network for Mood and Anxiety Treatments (CANMAT) clinical guidelines for the management of major depressive disorder in adults. II. Psychotherapy alone or in combination with antidepressant medication. *J Affect Disord* 2009; 117(Suppl 1):S15–S25
- World Health Organization: *mhGAP Intervention Guide for Mental, Neurological and Substance Use Disorders in Non-Specialized Health Settings: Mental Health Gap Action Programme (mhGAP)*. Geneva: World Health Organization, 2010.
- Australian and New Zealand clinical practice guidelines for the treatment of depression. Royal Australian and New Zealand College of Psychiatrists Clinical Practice Guidelines Team for Depression; 2004 [cited 2014 Feb 24]; Available from: https://www.ranzcp.org/Files/ranzcp-attachments/Resources/Publications/CPG/Clinician/CPG_Clinician_Full_Depression-pdf.aspx.
- Yager J, Devlin MJ, Halmi KA, Herzog DB, Mitchell JE III, Powers P, Zerbe KJ: *Practice Guideline for the Treatment of Patients with Eating Disorders*, 3rd ed. Arlington, VA, American Psychiatric Association, Inc, 2006
- Hirschfeld RMA, Bowden CL, Gitlin MJ, Keck PE, Suppes T, Thase ME, Perlis RH: *Practice Guideline for the Treatment of Patients with Bipolar Disorder*, 2nd ed. Arlington, VA, American Psychiatric Association, Inc, 2002
- Yatham LN, Kennedy SH, Parikh SV, Schaffer A, Beaulieu S, Alda M, O’Donovan C, Macqueen G, McIntyre RS, Sharma V, Ravindran A, Young LT, Milev R, Bond DJ, Frey BN, Goldstein BI, Lafer B, Birmaher B, Ha K, Nolen WA, Berk M: Canadian Network for Mood and Anxiety Treatments (CANMAT) and International Society for Bipolar Disorders (ISBD) collaborative update of CANMAT guidelines for the management of patients with bipolar disorder: update 2013. *Bipolar Disord* 2013; 15:1–44
- Australian and New Zealand clinical practice guidelines for the treatment of bipolar disorder. Royal Australian and New Zealand College of Psychiatrists Clinical Practice Guidelines Team for Bipolar Disorder; 2004 [cited 2014 Feb 24]; Available from: http://www.ranzcp.org/Files/ranzcp-attachments/Resources/Publications/CPG/Clinician/CPG_Clinician_Full_Bipolar-pdf.aspx.
- Segal ZV, Whitney DK, Lam RW; CANMAT Depression Work Group: Clinical guidelines for the treatment of depressive disorders. III. Psychotherapy. *Can J Psychiatry* 2001; 46(Suppl 1):29S–37S
- Bowlby J: *Attachment and loss*, vol. I: Attachment. New York, Basic Books, 1969
- Bowlby J: *Attachment and loss*, vol. 3: Loss: Sadness and Depression. New York, Basic Books, 1980
- Browne G, Harris T: *Social Origins of Depression: A Study of Psychiatric Disorders in Women*. New York, Free Press, 1978
- Meyer A: *Psychobiology; A Science of Man*. Oxford, UK, Charles C Thomas, 1957
- Sullivan HS: *The Interpersonal Theory of Psychiatry*. New York, WW Norton, 1953
- Holt-Lunstad J, Smith TB, Layton JB: Social relationships and mortality risk: a meta-analytic review. *PLoS Med* 2010; 7:e1000316
- Eisenberger NI, Cole SW: Social neuroscience and health: neurophysiological mechanisms linking social ties with physical health. *Nat Neurosci* 2012; 15:669–674
- Uchino BN: Social support and health: a review of physiological processes potentially underlying links to disease outcomes. *J Behav Med* 2006; 29: 377–387
- Cohen S: Social relationships and health. *Am Psychol* 2004; 59:676–684
- Ravitz P, Watson P, Grigoriadis S: *Interpersonal Psychotherapy for Depression*. Edited by Maunder R, Ravitz P. New York, WW Norton, 2013
- Lewinsohn PM, Rohde P, Seeley JR: Major depressive disorder in older adolescents: prevalence, risk factors, and clinical implications. *Clin Psychol Rev* 1998; 18:765–794
- Mufson L, Moreau D, Weissman MM, Wickramaratne P, Martin J, Samoilov A: Modification of interpersonal psychotherapy with depressed adolescents (IPT-A): phase I and II studies. *J Am Acad Child Adolesc Psychiatry* 1994; 33: 695–705
- Mufson L, Weissman MM, Moreau D, Garfinkel R: Efficacy of interpersonal psychotherapy for depressed adolescents. *Arch Gen Psychiatry* 1999; 56: 573–579
- Young JF, Mufson L, Davies M: Efficacy of interpersonal psychotherapy—adolescent skills training: an indicated preventive intervention for depression. *J Child Psychol Psychiatry* 2006; 47:1254–1262
- Moreau D, Mufson L, Weissman MM, Klerman GL: Interpersonal psychotherapy for adolescent depression: description of modification and preliminary application. *J Am Acad Child Adolesc Psychiatry* 1991; 30: 643–651
- Ehlers CL, Frank E, Kupfer DJ: Social zeitgebers and biological rhythms; a unified approach to understanding the etiology of depression. *Arch Gen Psychiatry* 1988; 45:948–952
- Ehlers CL, Kupfer DJ, Frank E, Monk TH: Biological rhythms and depression: The role of zeitgebers and zeitstorsers. *Depression* 1993; 1:285–293
- Monk TH, Flaherty JF, Frank E, Hoskinson K, Kupfer DJ: The Social rhythm metric; an instrument to quantify the daily rhythms of life. *J Nerv Ment Dis* 1990; 178:120–126
- Frank E, Kupfer DJ, Thase ME, Mallinger AG, Swartz HA, Fagiolini AM, Grochocinski V, Houck P, Scott J, Thompson W, Monk T: Two-year outcomes for interpersonal and social rhythm therapy in individuals with bipolar I disorder. *Arch Gen Psychiatry* 2005; 62:996–1004

44. Swartz HA, Levenson JC, Frank E: Psychotherapy for bipolar II disorder: The role of interpersonal and social rhythm therapy. *Professional Psychology: Research and Practice* 2012; 43:145–153
45. Frank E: Interpersonal and social rhythm therapy: a means of improving depression and preventing relapse in bipolar disorder. *J Clin Psychol* 2007; 63:463–473
46. Rucci P, Frank E, Kostelnik B, Fagiolini A, Mallinger AG, Swartz HA, Thase ME, Siegel L, Wilson D, Kupfer DJ: Suicide attempts in patients with bipolar I disorder during acute and maintenance phases of intensive treatment with pharmacotherapy and adjunctive psychotherapy. *Am J Psychiatry* 2002; 159:1160–1164
47. Hudson JI, Hiripi E, Pope HG Jr, Kessler RC: The prevalence and correlates of eating disorders in the national comorbidity survey replication. *Biol Psychiatry* 2007; 61:348–358
48. Arcelus J, Mitchell AJ, Wales J, Nielsen S: Mortality rates in patients with anorexia nervosa and other eating disorders; a meta-analysis of 36 studies. *Arch Gen Psychiatry* 2011; 68:724–731
49. Wilfley DE, Welch RR, Stein RI, Spurrell EB, Cohen LR, Saelens BE, Douchis JZ, Frank MA, Wiseman CV, Matt GE: A randomized comparison of group cognitive-behavioral therapy and group interpersonal psychotherapy for the treatment of overweight individuals with binge-eating disorder. *Arch Gen Psychiatry* 2002; 59:713–721
50. Carter CA, Jordan J, McIntosh VWW, Luty SE, McKenzie JM, Frampton CMA, Bulik CM, Joyce PR: The long-term efficacy of three psychotherapies for anorexia nervosa: a randomized, controlled trial. *Int J Eat Disord* 2011; 44:647–654
51. Demyttenaere K, Bruffaerts R, Posada-Villa J, Gasquet I, Kovess V, Lepine JP, Angermeyer MC, Bernert S, de Girolamo G, Morosini P, Polidori G, Kikkawa T, Kawakami N, Ono Y, Takeshima T, Uda H, Karam EG, Fayyad JA, Karam AN, Mneimneh ZN, Medina-Mora ME, Borges G, Lara C, de Graaf R, Ormel J, Gureje O, Shen Y, Huang Y, Zhang M, Alonso J, Haro JM, Vilagut G, Bromet EJ, Gluzman S, Webb C, Kessler RC, Merikangas KR, Anthony JC, Von Korff MR, Wang PS, Brugha TS, Aguilar-Gaxiola S, Lee S, Heeringa S, Pennell BE, Zaslavsky AM, Ustun TB, Chatterji S; WHO World Mental Health Survey Consortium: Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. *JAMA* 2004; 291:2581–2590
52. Bass J, Neugebauer R, Clougherty KF, Verdelli H, Wickramaratne P, Ndogoni L, Speelman L, Weissman M, Bolton P: Group interpersonal psychotherapy for depression in rural Uganda: 6-month outcomes: randomised controlled trial. *Br J Psychiatry* 2006; 188:567–573
53. Bolton P, Bass J, Betancourt T, Speelman L, Onyango G, Clougherty KF, Neugebauer R, Murray L, Verdelli H: Interventions for depression symptoms among adolescent survivors of war and displacement in northern Uganda: a randomized controlled trial. *JAMA* 2007; 298:519–527
54. Bolton P, Bass J, Neugebauer R, Verdelli H, Clougherty KF, Wickramaratne P, Speelman L, Ndogoni L, Weissman M: Group interpersonal psychotherapy for depression in rural Uganda: a randomized controlled trial. *JAMA* 2003; 289:3117–3124
55. Pain C, Wondimagegn D, Alem A, Ravitz P, Frank E, Nelson S, Wilson L, Baheretibeb Y, Bender A, Fekadu A, Chowdhary N, Hanlon C, Philpott J, Shibre T, Verdelli H: The Baber Project—Scaling up Interpersonal Psychotherapy (IPT) for Common Mental Disorders in Ethiopia. *Ethiopia, Grand Challenges*, 2013
56. Patel VH, Kirkwood BR, Pednekar S, Araya R, King M, Chisholm D, Simon G, Weiss H: Improving the outcomes of primary care attenders with common mental disorders in developing countries: a cluster randomized controlled trial of a collaborative stepped care intervention in Goa, India. *Trials* 2008; 9:4
57. Ravitz P, Wondimagegn D, Pain C, Alem A, Baheretibeb Y, Hanlon CFA, Fekadu A, Maunder R, Park J, Fefergrad M, Leszcz M: Psychotherapy knowledge translation: a review of education practices to transform mental health care, illustrated in Interpersonal Psychotherapy (IPT) and training applications in Canada and Ethiopia. *Am J Psychother* (in press)
58. Verdelli H, Clougherty K, Bolton P, Speelman L, Lincoln N, Bass J, Neugebauer R, Weissman MM: Adapting group interpersonal psychotherapy for a developing country: experience in rural Uganda. *World Psychiatry* 2003; 2:114–120
59. Grote NK, Swartz HA, Geibel SL, Zuckoff A, Houck PR, Frank E: A randomized controlled trial of culturally relevant, brief interpersonal psychotherapy for perinatal depression. *Psychiatr Serv* 2009; 60:313–321
60. Markowitz JC, Patel SR, Balan IC, Bell MA, Blanco C, Yellow Horse Brave Heart M, Sosa SB, Lewis-Fernández R: Toward an adaptation of interpersonal psychotherapy for Hispanic patients with DSM-IV major depressive disorder. *J Clin Psychiatry* 2009; 70:214–222
61. Betancourt TS, Bass J, Borisova I, Neugebauer R, Speelman L, Onyango G, Bolton P: Assessing local instrument reliability and validity: a field-based example from northern Uganda. *Soc Psychiatry Psychiatr Epidemiol* 2009; 44:685–692
62. Bolton P, Tang AM: An alternative approach to cross-cultural function assessment. *Soc Psychiatry Psychiatr Epidemiol* 2002; 37:537–543
63. Wilfley DE, MacKenzie KR, Welch RR, Ayres VE, Weissman M: *Interpersonal Psychotherapy for Group*. New York, Basic Books, 2000
64. Weissman M, Verdelli H: Interpersonal psychotherapy: evaluation, support, triage. *Clin Psychol Psychother* 2012; 19:106–112
65. Judd F, Weissman M, Davis J, Hodgins G, Piterman L: Interpersonal counselling in general practice. *Aust Fam Physician* 2004; 33:332–337
66. Kleinman A, Eisenberg L, Good B: Culture, illness, and care: clinical lessons from anthropologic and cross-cultural research. *Ann Intern Med* 1978; 88:251–258
67. Kirmayer LJ: Cultural variations in the clinical presentation of depression and anxiety: implications for diagnosis and treatment. *J Clin Psychiatry* 2001; 62(Suppl 13):22–28, discussion 29–30
68. Levenson JC, Frank E, Cheng Y, Rucci P, Janney CA, Houck P, Forgiome RN, Swartz HA, Cyranowski JM, Fagiolini A: Comparative outcomes among the problem areas of interpersonal psychotherapy for depression. *Depress Anxiety* 2010; 27:434–440
69. Pain C, Leszcz M, Hunter J, Maunder R, Ravitz P: *Psychotherapy Essentials to Go: Achieving Psychotherapy Effectiveness*. Edited by Maunder R, Ravitz P. New York, WW Norton, 2014
70. Allen JG, Fonagy P, Bateman AW: *Mentalizing in clinical practice*. Arlington, VA, American Psychiatric Association, 2008
71. Allen JG: *Restoring Mentalizing in Attachment Relationships: Treating Trauma with Plain Old Therapy*. Arlington, VA, American Psychiatric Association, 2013
72. Ravitz P, Maunder R, McBride C, (ed): *Improving IPT effectiveness—learning from treatment failures*. 5th International ISIPT Conference; 2013 Iowa City, Iowa.
73. Kawachi I, Berkman LF: Social ties and mental health. *J Urban Health* 2001; 78:458–467
74. Roth A, Fonagy P: *What Works for Whom: A Critical Review of Psychotherapy Research*, 2nd ed. New York, Guilford Press, 2005
75. Norcross JC (ed) *Psychotherapy Relationships That Work: Evidence-Based Responsiveness*, 2nd ed. New York, Oxford University Press, 2011
76. Constantino MJ, Manber R, DeGeorge J, McBride C, Ravitz P, Zuroff DC, Klein DN, Markowitz JC, Rothbaum BO, Thase ME, Arnow BA: Interpersonal styles of chronically depressed outpatients: profiles and therapeutic change. *Psychotherapy (Chic)* 2008; 45:491–506
77. Cyranowski JM, Bookwala J, Feske U, Houck P, Pilkonis P, Kostelnik B, Frank E: Adult attachment profiles, interpersonal difficulties, and response to interpersonal psychotherapy in women with recurrent major depression. *J Soc Clin Psychol* 2002; 21:191–217 doi: 10.1521/jscp.21.2.191.22514
78. Joyce PR, McKenzie JM, Carter JD, Rae AM, Luty SE, Frampton CM, Mulder RT: Temperament, character and personality disorders as predictors of response to interpersonal psychotherapy and cognitive-behavioural therapy for depression. *Br J Psychiatr* 2007; 190:503–508.
79. Kiesler DJ: *Contemporary Interpersonal Theory and Research: Personality, Psychopathology, and Psychotherapy*. New York, John Wiley & Sons, 1996
80. McBride C, Atkinson L, Quilty LC, Bagby RM: Attachment as moderator of treatment outcome in major depression: a randomized control trial of interpersonal psychotherapy versus cognitive behavior therapy. *J Consult Clin Psychol* 2006; 74:1041–1054
81. McBride C, Zuroff DC, Ravitz P, Koestner R, Moskowitz DS, Quilty L, Bagby RM: Autonomous and controlled motivation and interpersonal therapy for depression: moderating role of recurrent depression. *Br J Psychiatr* 2010; 49:529–545.
82. Ravitz P, Maunder R, McBride C: Attachment, contemporary interpersonal theory and IPT: an integration of theoretical, clinical, and empirical perspectives. *J Contemp Psychother* 2008; 38:11–21
83. Carter JD, Luty SE, McKenzie JM, Mulder RT, Frampton CM, Joyce PR: Patient predictors of response to cognitive behaviour therapy and interpersonal psychotherapy in a randomised clinical trial for depression. *J Affect Disord* 2011; 128:252–261
84. Ravitz P, McBride C, Maunder R: Failures in interpersonal psychotherapy (IPT): factors related to treatment resistances. *J Clin Psychol* 2011; 67:1129–1139
85. Norcross JC (ed): *Psychotherapy Relationships that Work: Evidence-Based Responsiveness*, 2nd ed. New York, Oxford University Press, 2011
86. Beck AT: *Cognitive Therapy of Depression*, New York, Guilford Press, 1979
87. Usher SF: *Introduction to Psychodynamic Psychotherapy Technique* 2nd ed.: New York, Routledge/Taylor & Francis Group, 2013.
88. Markowitz JC, Svartberg M, Swartz HA: Is IPT time-limited psychodynamic psychotherapy? *J Psychother Pract Res* 1998; 7:185–195
89. Hellerstein DJ, Pinsky H, Rosenthal RN, Klee S: Supportive therapy as the treatment model of choice. *J Psychother Pract Res* 1994; 3:300–306